

FAUNA AND PALAEOECOLOGY OF GRAN DOLINA-TD6 (SIERRA DE ATAPUERCA, SPAIN)

Jesús Rodríguez, Jan van der Made, Antonio Rosas & José María Bermúdez de Castro

The Sierra de Atapuerca is a low limestone range situated in central northern Spain, 15 Km east of Burgos city. Several karst cavities in the range are filled with Pleistocene sediments. Gran Dolina, one of these deposits, contains over 18 m of sediments, which can be divided into 11 lithostratigraphic levels named TD1-TD11, from bottom to top. A paleomagnetic reversal has been detected within the TD7 level of Gran Dolina (Parés & Pérez González, 1995) indicating a late Early Pleistocene age for TD1 to TD6 levels, and a Middle Pleistocene age for the upper levels. TD6 includes the Aurora stratum, where several stone tools and *Homo* antecedent remains were found (Carbonell *et al.*, 1995). The ungulate fauna from TD6 includes *Stephanorhinus etruscus*, a stenonine *Equus*, *Sus scrofa*, *Dama dama vallonensis*, *Cervus elaphus*, *Megaloceros* and *sp.*, *Bison sp.* This fauna is in agreement with a late early Pleistocene Age for this level. The microfauna from TDW4 and TDW4b is characteristic of the Biharian (Cuenca *et al.*, 1995).

The paleoecology of the lower levels of Gran Dolina has been studied using ecological diversity analysis (Andrews *et al.*, 1979). A data set of 92 recent mammalian faunas is used for comparison. The 5 assemblages studied (TDW4, TDW4b, TD6inf, TD6Au and TD8) share some characteristics that set them apart from all the recent communities analysed, namely a high proportion of large herbivores combined with the absence of arboreal and climbing mammals, and a low number of small carnivores. The abundance of large herbivores and the absence of arboreal mammals are considered unique features of the Pleistocene communities. The community structure of TD6inf, TD6Au, and TD8 resembles that of the recent European forest communities, however TDW4 and TDW4b have a community structure different from any recent community from the holartic region.

References

- ANDREWS P., LORD J.M. & NESBIT EVANS E.M., 1979. Patterns of ecological diversity in fossil and modern mammalian faunas. *Biol. Journ. Linn. Soc.* 11:177-205.
- CARBONELL E., BERMÚDEZ DE CASTRO J.M., ARSUAGA J.L., Díez J.C., ROSAS A., CUENCA-BESCÓS G., SALA R., MOSQUERA M., y RODRÍGUEZ X.P., 1995. Lower Pleistocene Hominids and Artefacts from Atapuerca-TD6 (Spain). *Science* 269:826-830.
- CUENCA G., CANUDO J.I. & LAPLANA C., 1995. Los arviclidos (Rodentia, Mammalia) de los niveles inferiores de Gran Dolina (Pleistoceno Inferior, Atapuerca, Burgos, España). *Rev. Esp. Pal.*, 10(2):202-218.
- PARES J.M. & PEREZ GONZALEZ A., 1995. Paleomagnetic age for hominid fossils at Atapuerca archaeological site, Spain. *Science* 269:830-832.